

This article is written by students. It may include omissions and imperfections,  
as far as possible reported by our reviewers in the editorial notes

# Are all infinities the same?

Year 2023–2024

George Aftene Bogdan, Albert Matei, Alexandra Burbulea, Andrei Razvan Damoc,  
Alexandra Lapusneanu, Mihai Maxim, Tudor Stefan Pascari, students in 9<sup>th</sup> grade

**School:** Colegiul Național C. *Negruzzi*, Iași

**Teachers:** Adrian Zanoschi, Colegiul Național C. *Negruzzi* and Razvan-Andrei Morariu, student at University of Vienna

**Researcher:** Razvan-Andrei Morariu, student at University of Vienna

## 1. PRESENTATION OF THE RESEARCH TOPIC

The concept of infinity has long captivated the minds of mathematicians, leading to profound insights and paradoxes. As we embark on this exploration, we will venture beyond the familiar realms of finite numbers and immerse ourselves in the abstract landscape where infinite sets reside. Our focus will be on understanding the diverse sizes of infinities and the methods used to compare them.


## 2. BRIEF PRESENTATION OF THE RESULTS OBTAINED

Before stating the problem, we will present the definition of two sets having the same cardinality.

**Definition 1:** Two sets  $A$  and  $B$  are considered to have the same cardinality (i.e.  $|A| = |B|$ ) if there is a bijective function between set  $A$  and set. [\[1\]](#)

Infinity is a really important and interesting concept in mathematics. Even though it is an abstract concept, we have to think about it right from the time we start learning about natural numbers because there are just endless natural numbers. After that, we can also think about there being endless integers or even endless prime numbers. However, even if infinity might sound like the biggest way to describe something, are all infinities as big as each other? Can one infinite set be as big as another infinite set even if one includes strictly the other? Is it possible to find different types of infinities? To address these curiosities and understand this concept better, we will try to answer the following questions:

1. Is the set of prime numbers infinite?
2. Do the set of natural numbers have the same cardinality as the set of prime numbers?
3. Can we compare the cardinality of  $\mathbb{N}$  with the cardinality of  $\mathbb{Z}$ ?
4. Can we compare the cardinality of  $\mathbb{N}$  and the cardinality of  $\mathbb{Q}$ ?
5. Are there sets with a higher cardinality than  $\mathbb{Q}$ ?
6. Does  $[0, 1]$  have the same cardinality as any other closed interval  $[a, b]$ ?
7. Is it also true that the interval  $(0, 1)$  has the same cardinality as any other open interval  $(a, b)$ ?
8. Does  $[0, 1]$  have the same cardinality as  $\mathbb{R}$ ?



### 3. SOLUTION

**Definition 2:** A set  $A$  is called countable if there is an injection  $f: A \rightarrow \mathbb{N}$ .

**Remark 1:** Any finite set  $A$  is countable because one can consider  $A = \{a_1, a_2, \dots, a_n\}$  and let  $f: A \rightarrow \mathbb{N}$ ,  $f(a_i) = i$ , for  $i \in \{1, 2, \dots, |A|\}$  an injection from  $A$  into  $\mathbb{N}$ . [2]

**Remark 2:** In literature a set of this type is called countable because one can make a list out of its elements (i.e. assign a numbered position for every element in the set).

**Observation:** We will consider the following theorems to be true:

**Theorem 1:** If there is an injective function from  $A$  into  $B$  and also an injective function from  $B$  into  $A$ , then  $A$  and  $B$  have the same cardinality. [3]

**Theorem 2:** If there is no surjection between  $A$  and  $B$ , this means that  $|A| < |B|$ . [4]

**Theorem 3:** If  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ .

**Remark 3:** There is an important intuition of **Definition 1**. Since a bijection assigns every element from a set to a unique distinct element from another set and covers both sets completely, it means that intuitively the two sets have the exact same size.

**Remark 4:** There is an intuitive relationship between **Theorem 1** and **Theorem 3**. Creating an injection from  $A$  to  $B$  proves intuitively that  $A$  and  $B$  because we can assign any element in  $A$  to a distinct element from  $B$ .

**Remark 5:** There is an intuitive explanation of **Theorem 2**. If there is no surjection between  $A$  and  $B$ , it means that one cannot cover the entire set  $B$  with the elements of  $A$ , so  $|A| < |B|$ .

### QUESTION 1: Is the set of prime numbers infinite?

Our intuition was that the set of prime numbers is infinite. To show this, we supposed, for the sake of contradiction, that there are only finitely many prime numbers, which can be listed as  $\{p_1, p_2, p_3, \dots, p_n\}$ .

Now, let's construct a new number, which is one more than the product of all these primes:  $N = p_1 \cdot p_2 \cdot p_3 \cdot \dots \cdot p_n + 1$ .

$N$  is clearly greater than 1 because it is one more than the product of prime numbers. So,  $N$  must be either composite or prime.

**We will show that  $N$  cannot be composite:**

If  $N$  is composite, it should have at least one prime divisor strictly smaller than itself. However, none of the primes  $p_1, p_2, p_3, \dots, p_n$  can be a divisor of  $N$ , because when  $N$  is divided by any of them, there is a remainder of 1. Therefore,  $N$  must have a prime factor that is not in our original list. In consequence, we reach a contradiction which means that our assumption that the set of prime numbers is finite is false. Therefore, there are indeed an infinite number of prime numbers. [5]

### QUESTION 2: Do the set of natural numbers have the same cardinality as the set of prime numbers?

We will denote by  $P$  the set of prime numbers which we already now is infinite.  $P = \{p_1, p_2, p_3, \dots, p_n, \dots\}$ , where  $p_1 = 2 < p_2 = 3 < p_3 = 5 \dots$ . [6]

We will consider the function  $f: \mathbb{N} \rightarrow P$ , where  $f(n) = p_{n+1}$ .

We will prove that  $f$  is bijective in 2 steps:

**I.** Firstly, we prove that  $f$  is **injective**.

Let us consider two natural numbers  $a$  and  $b$  such that  $f(a) = f(b)$ .

$f(a) = f(b)$  means that  $p_{a+1} = p_{b+1}$ . Since we considered the prime numbers to be arranged in strictly increasing order this means that  $a = b$ .

**II.** Secondly, we must prove that  $f$  is **surjective**.

To prove this, we must show that for any prime number  $p \in P$  there is at least one natural number  $x$  such that  $f(x) = p$ . Knowing that  $p$  is prime, there exists a  $n \in \mathbb{N}$  such that  $p$  is the  $n + 1$ -th prime number in the list of prime numbers arranged in increasing order. So, there is  $x = n$  such that  $f(x) = p$ . This means that the function is surjective.

In conclusion, because the function is both injective and surjective,  $f(n) = p_{n+1}$  is a bijection between  $\mathbb{N}$  and  $P$ . So, we can conclude that  $|\mathbb{N}| = |P|$ .

### QUESTION 3: Can we compare the cardinality of $\mathbb{N}$ with the cardinality of $\mathbb{Z}$ ?

We will try to find a bijective function between  $\mathbb{N}$  and  $\mathbb{Z}$ .

Let's consider the function  $f: \mathbb{N} \rightarrow \mathbb{Z}$  defined by

1.  $f(x) = -\frac{x}{2}$  for any even  $x$ .
2.  $f(x) = \frac{x+1}{2}$  for any odd  $x$ .

We want to prove that function  $f$  is bijective. We will prove this in 2 steps, by proving that  $f$  is injective and surjective.

Function  $f: \mathbb{N} \rightarrow \mathbb{Z}$  is injective if  $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$  for any  $x_1, x_2 \in \mathbb{N}$ .

First of all, one can observe that  $f$  is strictly positive for odd numbers and negative for even numbers. So, we can have  $f(x) = f(y)$  only if  $2|(x - y)$ .

If  $x_1$  and  $x_2$  are even,  $f(x_1) = f(x_2) \Leftrightarrow -\frac{x_1}{2} = -\frac{x_2}{2} \Leftrightarrow x_1 = x_2$ .

If  $x_1$  and  $x_2$  are odd,  $f(x_1) = f(x_2) \Leftrightarrow \frac{x_1+1}{2} = \frac{x_2+1}{2} \Leftrightarrow x_1 = x_2$ .

**So,  $f$  is injective.**

Now we move to surjectivity.

Function  $f: \mathbb{N} \rightarrow \mathbb{Z}$  is surjective if, for any  $y \in \mathbb{Z}$ , there is an  $x \in \mathbb{N}$  such that  $y = f(x)$ .

Any  $y > 0$  can be expressed as  $\frac{x+1}{2}$  by taking  $x = 2y - 1$ .

Any  $y \leq 0$  can be expressed as  $-\frac{x}{2}$  by taking  $x = -2y$ .

**This shows that  $f$  is surjective.**

Function  $f$  is both injective and surjective, **so  $f$  is bijective**. In conclusion, natural numbers and integers have the same cardinality.

**So, from now on, we can use the cardinality of  $\mathbb{N}$  and  $\mathbb{Z}$  interchangeably.**

**Remark:** The cardinality of any infinite countable set is denoted by  $\aleph_0$  [7].

#### **QUESTION 4: Can we compare the cardinality of $\mathbb{N}$ with the cardinality of $\mathbb{Q}$ ?**

We will try to find one injective function from  $\mathbb{N}$  to  $\mathbb{Q}$ , and one injective function from  $\mathbb{Q}$  to  $\mathbb{N}$ .

The function from  $\mathbb{N}$  to  $\mathbb{Q}$  is natural:  $g: \mathbb{N} \rightarrow \mathbb{Q}$ , where  $g(x) = x$ . This function is injective and well defined. We will mark this finding as **(1)**.

Now we will focus on the other injection. Since every rational number  $r$  can be written as  $\frac{a}{b}$ , where  $(a, b) = 1$  and  $b > 0$ . Let  $f: \mathbb{Q} \rightarrow \mathbb{N}$  be defined as

$$1) f(r) = 2^a \cdot 3^b \text{ for } r = \frac{a}{b}, \text{ if } a > 0$$

$$2) f(r) = 5^{-a} \cdot 3^b \text{ for } r = \frac{a}{b}, \text{ if } a \leq 0.$$

One can show that that  $\text{Im } f \subset \mathbb{N}$ , because  $2^a \in \mathbb{N}$  when  $a > 0$ ,  $5^{-a} \in \mathbb{N}$  when  $a \leq 0$ , and  $3^b \in \mathbb{N}$  when  $b > 0$ . So,  $f$  is well defined. Now, we will prove that  $f$  is injective. Firstly, if  $f(x) = f(y)$ ,  $(x, y)$  must have the same sign.

So, there are two cases given their sign.

If  $f(x) = 2^a \cdot 3^b$ ,  $f(y) = 2^c \cdot 3^d$  and  $f(x) = f(y)$  it will result that the powers of 2 will be equal (i.e.  $a = c$ ) and the powers of 3 will be also equal (i.e.  $b = d$ ). In a similar way, if we consider that  $f(x) = 5^{-a} \cdot 3^b$ ,  $f(y) = 5^{-c} \cdot 3^d$  and  $f(x) = f(y)$  it will result that the powers of 5 will be equal (i.e.  $a = c$ ) and the powers of 3 will be also equal (i.e.  $b = d$ ). That shows the function  $f$  is injective because  $f(x) = f(y) \Leftrightarrow x = y$ .

Therefore, the function  $f$  is **well defined and injective**. We will mark this finding as **(2)**.

Combining **(1)** and **(2)**, with the help of **Theorem 1**, we will obtain the following important result:  $|\mathbb{N}| = |\mathbb{Q}|$ .

### QUESTION 5: Are there sets with a higher cardinality than $\mathbb{Q}$ ?

We will prove that the cardinality of  $[0, 1]$  is strictly higher than the cardinality of  $\mathbb{Q}$ .

To this end, we will prove that there is no surjective function between natural numbers and the interval  $[0, 1]$ . This will show that the cardinality of  $[0, 1]$  is higher than the cardinality of  $\mathbb{Q}$  because we already showed that  $|\mathbb{N}| = |\mathbb{Q}|$ .

Firstly, we will suppose that such a bijection exists. Then, using Cantor's Diagonalization, we will obtain a list as follows: [8]

0 ← → 0.52149356 ...  
1 ← → 0.14162985 ...  
2 ← → 0.94782712 ...  
3 ← → 0.53098175 ...  
⋮

The number circled in the diagonal is some real number  $r$ , since it is an infinite decimal expansion. Now consider the real number  $s$  obtained by modifying every digit of  $r$ , say by replacing each digit that is not 6 with 6 and each 6 with 7. We claim that  $s$  does not occur in our infinite list of real numbers. Suppose by sake of contradiction that it does, and that it is the  $n$ -th number in the list. Then  $r$  and  $s$  differ in the  $n$ -th digit due to the construction of  $s$ . Moreover, since the decimals of  $s$  can be 6 or 7 only, it has a unique decimal expansion, which means that it cannot appear in a different form in the list. So, we have a real number  $s$  that is not in the image of our supposed surjective function. [9]

This **contradicts the assertion** that  $f$  is a surjection. Thus, there is no surjection between the natural numbers and the interval  $[0, 1]$ . So, the cardinality of  $[0, 1]$  is **strictly higher** than the cardinality of  $\mathbb{N}$  due to **Theorem 2**.

**Observation:** The main ideas of this solution were taken from <https://inst.eecs.berkeley.edu/~cs70/sp07/lec/lecture27.pdf>.

### QUESTION 6: Does $[0, 1]$ have the same cardinality as any other closed interval $[a, b]$ ?

We want to establish a bijection between the intervals  $[a, b]$  and  $[0, 1]$  in order to prove that they have the same cardinality.

We will take the function  $f: [a, b] \rightarrow [0, 1]$  with  $f(x) = \frac{x-a}{b-a}$ . We can observe that  $f$  is well defined because  $\text{Im } f \subset [0, 1]$ . [10]

Firstly, we will prove that the function  $f$  is injective.

We can assume that  $f(x_1) = f(x_2)$  for two values  $x_1, x_2 \in [a, b]$ . One can observe that

$$f(x_1) = f(x_2) \Leftrightarrow \frac{x_1 - a}{b - a} = \frac{x_2 - a}{b - a} \Leftrightarrow x_1 - a = x_2 - a \Leftrightarrow x_1 = x_2.$$

So,  $f$  is **injective**. Now, we need to prove that  $f$  is surjective. For that, we have to prove that for every  $y \in [0, 1]$  there is a value  $x \in [a, b]$  such that  $f(x) = y$ . Since

$$f(x) = y \Leftrightarrow \frac{x - a}{b - a} = y \Leftrightarrow x - a = y \cdot (b - a).$$

In consequence, for every  $y \in [0, 1]$ , there exists  $x = y \cdot (b - a) + a \in [a, b]$  such that  $f(x) = y$ . This means that  $f$  is **surjective**.

If  $f$  is injective and surjective,  $f$  is **bijective**.

In consequence  $[0, 1]$  has the same cardinality as any other closed interval  $[a, b]$ .

### QUESTION 7: Is it also true that $(0, 1)$ has the same cardinality as any other open interval $(a, b)$ ?

Using the function, we already built at the previous point  $f: [a, b] \rightarrow [0, 1]$  with  $f(x) = \frac{x-a}{b-a}$ , we will prove that there is a bijective function  $F: (a, b) \rightarrow (0, 1)$ .

Knowing that  $f(a) = 0$  and  $f(b) = 1$  we will consider  $F$  to be the restriction of  $f$  on  $(a, b)$ . It is clear that  $F$  will be a bijection between  $(a, b)$  and  $(0, 1)$ .

**Remark:** Since  $\left[ a + \frac{b-a}{4}, b - \frac{b-a}{4} \right] \subset (a, b) \subset [a - 1, b + 1]$  and all closed intervals have the same cardinality, we can conclude that open intervals and closed intervals have the same cardinality due to Theorem 3.

### QUESTION 8: Does $[0, 1]$ have the same cardinality as $\mathbb{R}$ ?

We are interested in this because we showed that  $[0, 1]$  has a higher cardinality than  $\mathbb{Q}$ .

We want to show that  $[0, 1]$  has the same cardinality as  $\mathbb{R}$ . In order to do this we will find an injection  $f: [0, 1] \rightarrow \mathbb{R}$ , and an injection  $g: \mathbb{R} \rightarrow [0, 1]$ .

$f$  can be chosen as  $f(x) = x$ . This function is **well defined and injective**.

For  $g: \mathbb{R} \rightarrow [0, 1]$ , we propose to take  $g(x) = \frac{1}{1+2^x}$ . Now we have to prove 2 things: that  $g$  is well defined (i.e.  $\text{Im } g \subseteq [0, 1]$ ) and that  $g$  is injective.

1.  $\text{Im } g \subseteq [0, 1]$

It is clear that  $g(x) > 0, \forall x \in \mathbb{R}$  Also, since  $2^x > 0 \Rightarrow g(x) < \frac{1}{1} = 1, \forall x \in \mathbb{R}$ . So,  $0 < g(x) < 1, \forall x \in \mathbb{R}$ .

2.  $g$  is injective

Since  $g(x) = g(y) \Leftrightarrow \frac{1}{1+2^x} = \frac{1}{1+2^y} \Leftrightarrow 1+2^x = 1+2^y \Leftrightarrow 2^x = 2^y \Leftrightarrow x = y$ , we can conclude that  $g$  is injective.

In conclusion,  $g$  is **well defined and injective**.

Having found an injection from  $[0, 1]$  to  $\mathbb{R}$  and from  $\mathbb{R}$  to  $[0, 1]$  we can conclude that  $\mathbb{R}$  and  $[0, 1]$  have the same cardinality due to **Theorem 1**.

## 4. CONCLUSION

**In order to solve this problem, we have used injective, surjective and bijective functions to compare the cardinality of different sets. The central theorems that we used are enumerated in the beginning of our solution.**

In the first part we have obtained a set of results proving that there is an infinite number of prime numbers and that the set of prime numbers has the same cardinality as the set of natural numbers, integers, and rational numbers. In the second part, we managed to prove that there are infinities that have a higher cardinality than the set of rational numbers. We found that the cardinality of any closed or open interval is bigger than  $\aleph_0$  and is equal to the cardinality of  $\mathbb{R}$ .

## EDITION NOTES

[1] Given that this is a definition, it sounds inappropriate to say: "...  $A$  and  $B$  are considered to have the same cardinality ...". It would be preferable: "We say that  $A$  and  $B$  have the same cardinality (and we write  $|A| = |B|$ ) ..."

[2] The symbol  $|A|$  has not been defined. A meaning has only been given to equalities of the form  $|A| = |B|$ . It should have been stated explicitly that, if  $A$  is a finite set,  $|A|$  denotes the number of its elements.

[3] This theorem is known as the *Cantor–Schroeder–Bernstein* theorem.

[4] The expression  $|A| < |B|$  has not been defined in the work. So, this so-called "theorem" seems more like a definition. A correct expression of the definition could be: We write  $|A| < |B|$  to mean that there is an injection, but no surjection, from  $|A|$  into  $|B|$ .

It must be noted that a correct definition of  $|A| < |B|$  also requires the existence of an injective function from  $A$  to  $B$ . It is not enough that no surjection from  $A$  to  $B$  exists.

One can then define  $|A| \leq |B|$  in the obvious way. It must be observed that, on the basis of this definition, the following Theorem 3 is just a rewriting of Theorem 1.

[5] This proof dates back to Euclid.

[6] If we write  $P = \{p_1, p_2, p_3, \dots, p_n, \dots\}$ , we are already assuming that there is a bijection between  $\mathbb{N}$  and  $P$ .

[7] Observe that only the comparison of cardinalities has been defined in the work. So, the expression "cardinality of a set" has no meaning in context of this paper (see also Note [2]).

In Set Theory the cardinality of a set is actually defined, but the definition involves some advanced notions.

[8] The list can be considered because we are assuming that the interval  $[0,1]$  (in  $\mathbb{R}$ ) has the same cardinality as  $\mathbb{N}$ . Cantor's Diagonalization is not used. It is used, instead, in the proof below the figure.

[9] The final part of this proof is a bit confused and can be simplified. It should be firstly emphasized that  $r$  and the  $n$ -th number in the list have the same  $n$ -th digit. So, the assumption that  $s$  is the  $n$ -th number in the list leads to a contradiction because, by construction, the  $n$ -th digit of  $s$  is different from that of  $r$ .

[10] This note is just a matter of definition. The symbol  $\subset$  generally denotes proper inclusion. Since the image of  $f$  is the whole interval  $[0,1]$ , the symbol  $\subseteq$  would be preferable.